

A. diagnose de Arabis Lusitânica Boiss.
é a seguinte: (Sect. Monanthemum)

A. perennis vel biennis caule erecto folioso
simplici pilis adpressissimis ramosis vel centro
affixis simplicibus hirsuto superne glabrescenti.
foliis pube ramosa adpersis oblongis obtuse
denticulatis obtusis radicalibus in petiolum
attenuatis caulinis sessilibus basi truncato-
auriculatis, racemo florum terminali per an-
thesin denso dein elongato, siliguis strictissimis
longis linearibus planissimis nitidis, torulosis,
valvularum nervis marginalibus validis in-
ter medio nervis parallelis stipato, seminibus
angustissime marginatis.

Hab. in collinis Castelemaedon prope Lacer (Hel.)
in agris in cultis prope Olysiporto nem.

Affinis habitu et multis characteribus *A. sagittatae*
et *A. hispanicae*. Scop. Haec differt indumento longio-
ri patenti, illa foliis caulinis basi evidenter cordatis,
florum racemis sub anthesi elongatis basiusculis,
utroque siliguis brevioribus angustioribus ob val-
vulas subcarinatas compresso-subtetragonis.

Siliguae *A. Lusitanicae* 1/2 pollices longae
lineae 3/4 latae.

Disponitur superiori de Franco postrema de quem é
De V. Ex

Lisboa 2/9/84

M. M. am. e m. abri, 84

António Ricardo de Caceres





